

12. Exercise sheet

Due-date: Friday, 11.07.2014, *before* the lecture has started

This is the second last exercise sheet of the second half of the semester.

Exercise 67

4 points

Let E be a finite set, $c : E \rightarrow \mathbb{R} \setminus \{0\}$ and $\mathcal{F} \subseteq 2^E$ such that (E, \mathcal{F}) is a matroid. Moreover, let $X_1, X_2 \in \mathcal{F}$ be two optimal solutions to the corresponding maximization problem, i. e.,

$$c(X_1) = c(X_2) = \max\{c(X) \mid X \in \mathcal{F}\},$$

and $e \in X_1$. Prove that there exists an $e' \in X_2$ with $c(e') = c(e)$. In particular, if $c(e_1) \neq c(e_2)$ for all $e_1 \neq e_2$, then there exists a unique optimal solution.

Exercise 68

4 points

Let $G = (V, E)$ be an undirected graph and

$$\mathcal{F} := \{F \subseteq V \mid \text{there is a matching } M \text{ of } G \text{ covering all nodes in } F\}.$$

Prove that (V, \mathcal{F}) is a matroid.

Exercise 69

4 points

Prove Theorem 16.8:

Let E be a finite set and $\mathcal{B} \subseteq 2^E$. Then, \mathcal{B} is the set of bases of some matroid (E, \mathcal{F}) if and only if

(a) $\mathcal{B} \neq \emptyset$;

(b) $B_1, B_2 \in \mathcal{B}, e \in B_1 \setminus B_2 \implies \exists f \in B_2 \setminus B_1$ with $(B_1 \setminus \{e\}) \cup \{f\} \in \mathcal{B}$.

Exercise 70


3 points

Prove that the rank function of a matroid is submodular.

Exercise 71

5 bonus points

Consider the matroid intersection problem on the graphic matroids defined by the following graphs:



For $X := \{a, b, c\}$, construct the digraph D_X . Find an S_X - T_X -path for which the corresponding augmentation does *not* yield a common independent set. Find a shortest S_X - T_X -path and construct an optimum solution to the matroid intersection problem.

Exercise 72**Tutorial session – 0 points**

Let E be a finite set, $c : E \rightarrow \mathbb{R}$ and $\mathcal{F} \subseteq 2^E$ such that (E, \mathcal{F}) is a matroid. Prove that the greedy algorithm yields a basis B of E maximizing $\sum_{e \in B} c(e)$.

Exercise 73**Tutorial session – 0 points**

Let $G = (V, E)$ be a bipartite graph and

$$\mathcal{F} := \{M \subseteq E \mid M \text{ is a matching in } G\} .$$

Prove that the independence system (E, \mathcal{F}) is the intersection of two matroids.

Exercise 74**Tutorial session – 0 points**

Let E be a finite set and $f : 2^E \rightarrow \mathbb{R}$. Prove that the following three statements are equivalent:

- (a) $f(A \cup B) + f(A \cap B) \leq f(A) + f(B)$ for all $A, B \subseteq E$;
- (b) $f(S \cup \{j\}) - f(S) \geq f(T \cup \{j\}) - f(T)$ for all $j \in E$ and $S \subseteq T \subseteq E \setminus \{j\}$;
- (c) $f(S \cup \{j\}) - f(S) \geq f(S \cup \{j, k\}) - f(S \cup \{k\})$ for all $j \neq k \in E$ and $S \subseteq E \setminus \{j, k\}$.

A function f with these properties is called *submodular*.